





MANAGEMENT GUIDE

PARENT STOCK PREMIUM MINI



1. KEY MESSAGES FOR REARING PERIOD (0–20 WEEKS): MANAGEMENT, EQUIPMENT AND STOCKING DENSITY	3
2. KEY MESSAGES FOR PRODUCTION PERIOD (20–65 WEEKS): MANAGEMENT, EQUIPMENT AND STOCKING DENSITY	5
3. REARING PERIOD (0–20 WEEKS)	6
HOW TO HAVE A SUCCESSFUL BROODING PERIOD?	6
HOW TO CHOOSE THE OPTIMUM LIGHTING PROGRAMME? DARK PRODUCTION HOUSE	7
HOW TO CHOOSE THE OPTIMUM LIGHTING PROGRAMME? OPEN-SIDED PRODUCTION HOUSE	8
HOW TO FOLLOW THE BODYWEIGHT TARGETS?	9
HOW TO REACH AND MAINTAIN GOOD FLOCK UNIFORMITY AND GOOD APPETITE?	10
HOW TO PROMOTE GOOD BEHAVIOUR AND PROVIDE GOOD ANIMAL WELFARE?	11
4. PRODUCTION PERIOD (20–65 WEEKS)	12
HOW TO MAKE THE FIRST 10 DAYS A SUCCESS? FOCUS ON LIGHT	12
HOW TO MAKE THE FIRST 10 DAYS A SUCCESS? FOCUS ON FEED PROGRAMME	13
HOW TO MAKE THE FIRST 10 DAYS A SUCCESS? FOCUS ON FEED DISTRIBUTION	14
HOW TO MAKE THE FIRST 10 DAYS A SUCCESS? FOCUS ON NEST LAYING	15
HOW TO MAXIMISE THE PERSISTENCY OF LAY?	16
5. BIOSECURITY	17

1. KEY MESSAGES FOR REARING PERIOD (0–20 WEEKS): MANAGEMENT, EQUIPMENT AND STOCKING DENSITY



1. KEY MESSAGES FOR REARING PERIOD (0–20 WEEKS): MANAGEMENT, EQUIPMENT AND STOCKING DENSITY

Successful rearing = more than 80% females within the flock receptive (mature but not too early) at the time of the light stimulation (148–154 days according to the female breed)

Flock management to be adapted according to the local rearing conditions (house type, climate...).

- Develop a good frame size by achieving bodyweight objectives at 4–6 weeks. *Ad libitum* feeding during the first 3 weeks and until 4 weeks of age if the 3 weeks bodyweight target is not achieved.

- Monitor sexual maturity to prevent the onset of lay occurring too early:

- By applying a slow step-down lighting programme during the first 10–12 weeks (to be adapted to the female breed and the local rearing conditions).
- By following the bodyweight objectives paying attention to weekly growth around 6–8 weeks and 16–22 weeks to ensure females are not too heavy.
- By avoiding disruption to growth and uniformity between 18–22 weeks, especially when doing transfer and doing vaccination.

- Provide 5 cm/bird of perch space or 4 m² of platform/1 000 birds from 28 days to train the birds to jump up to the nests.

- Develop a good appetite to prepare the onset of lay and a good flock uniformity above 85% (CV < 7.5). Maintain feed clean-up time less than 7 hours.

- All the life of the flock:

- Assure good distribution of both feed and water.
- Maintain good litter quality.
- Maintain proper environment and air quality for a healthy flock.
- Weigh 3–5% of the flock or at least 100 birds per pen per week.

i



Performance Objectives
JA57 Ki PS



Performance Objectives
JA87 PS



Performance Objectives
REDBRO MINI PS



Performance Objectives
P6N PS



Performance Objectives
JA57 PS



Nutrition
Recommendations

EQUIPMENT AND STOCKING DENSITY DURING REARING	Stocking density *	7–10 birds/m ² (< 7 birds/m ² in hot climate)
	Brooders	1 for 500 chicks
	Watering – round	1 for 80 birds
	Watering – nipple	1 for 8–10 birds
	Feeding – chain	10–12 cm/bird (5–6m/100 birds)
	Feeding pans – round	1 for 13–15 birds
	Feeding pans – oval	1 for 14–16 birds
	Spin feeder	1 for 1 500–1 800 birds
	Feed distribution time	< 4 minutes

* Stocking density may be subject to local regulations which must be observed at all times.



Dark rearing house – good perch access

2. KEY MESSAGES FOR PRODUCTION PERIOD (20–65 WEEKS): MANAGEMENT, EQUIPMENT AND STOCKING DENSITY

GROWTH: ACHIEVE 5 % PRODUCTION NOT BEFORE 22–23 WEEKS OF AGE (ACCORDING TO THE FEMALE BREED).

- Avoid disruption to growth and uniformity between 20–23 weeks, especially when changing feed type and mixing males.
- Appropriate eating behaviour between first egg and depletion. Check feed distribution on a regular basis. → See Poster *“Feed distribution”*.
- Ensure weekly growth rate is on target. Weigh 3–5% of the flock or least 100 birds per pen per week.

LIGHT

- From 148–154 days (according to the female breed), provide a first increase in daylength of at least 2 h combined with an increase of the light intensity (60–80 lux in dark houses) then + 1 h light / week until 14 h or 15h light in dark production house and 15 h light in open-sided production house.

- It is recommended not to exceed 16 h of light duration.

FEED: PROMOTE A GOOD EGG PRODUCTION FROM THE FIRST EGGS.

- Give daily feed increase from 5–10% daily production and reach the feed peak no later than 55–60% daily production.
- Control male access to female feeders by using the correct grill size or chain feeders (43–45 mm wide x 55 mm high). Consider adding a pipe of 20–25 mm width to the top of the grill if a taller grill is used and adjusting pan feeders to the minimum width and height setting.
- When egg weight exceeds 60 g it is recommended to use Breeder II feed.
- Plan feed reduction according to egg production, egg weight, female bodyweight and abdominal fat.

NEST EGGS: PREPARE THE FLOCK PROPERLY TO MAXIMISE NEST LAYING.

- Timely reaction to laying on the floor with corrective action.
 - Frequent collection of floor eggs from the first eggs remains a key practice.
 - The mating ratio will be defined according to the type of males being used and the level of sexual maturity at 20–22 weeks. → See Management guide *“PS males mated with Premium females”*.
- A detailed checklist is available from your local Hubbard Technical Specialist.

WATER: MAXIMISE THE FEMALE LIVEABILITY AND THE LITTER QUALITY.

- Control water quality: pH = 5.5 to 6.8
- Oxidation Reduction Potential (ORP) or Redox potential > 600 mV with chlorine
- Chlorine concentration at the end of the pipe = 1 ppm (Acceptable range: 0.5 to 3 ppm).



Open-sided laying house

EQUIPMENT AND STOCKING DENSITY DURING PRODUCTION	Stocking density	<7.5 females/available m ² (excl. egg belt and service room) < 6 birds/available m ² in hot climate
	Watering – round	1 for 80 females
	Watering – nipple	1 for 6 to 10 females (nipple flow 70–100 ml/min*)
	Feeding – chain	12.5–14 cm feeder space per female/ 6.25–7 m length for 100 females**
	Feeding pans – round ø 35 cm	1 for 13–15 females **
	Feeding pans – oval	1 for 14–16 females **
	Feed distribution time	< 4 minutes
	Nests	1 manual nest/4 females 80–90 females/linear meter of automatic nest
	Required ventilation	0.6 to 6 m ³ /kg liveweight/hour
	Air Speed	2.5 m/sec up to 3 m/sec in humid and warm climate (tunnel ventilation)
	Light intensity	60–80 lux

* Some nipple drinker systems are designed to operate with lower flow rates for breeders so check the manufacturer's recommendations or seek advice from your Hubbard Technical Specialist.

** Feeding space may be subject to local regulations which must be observed at all times.

[Male management guide](#)


[Nutrition Recommendations](#)

[Hubbard Poster Feed distribution](#)

[Hubbard Bulletin Water Quality](#)

3. REARING PERIOD (0-20 WEEKS): HOW TO HAVE A SUCCESSFUL BROODING PERIOD?


The key management practices are shown in our brooding poster. → See Poster **“Brooding PS”**. A close observation of the chicks during the first two weeks in their environment will make brooding successful (optimal growth and good feed intake) by adjusting light, feed, water, equipment, heat and minimum ventilation.



Optimal crumble starter feed

KEY POINTS TO PROMOTE EARLY WATER AND FEED INTAKE

- Choose an appropriate density (25-30 birds/m²) during the first 3 days. Chicks should never be away more than one meter from a drinker or a feeder.
- If permitted by local regulations, use an intermittent lighting programme from day 1 to day 28. Always respect the recommended hours of light below and a minimum of 4 cycles of light/dark per day. Ensure to be present in the house to observe the chick behaviour on the first few times the lights are switched on and off.
- Place chick paper with feed on at least 50%



Intermittent lighting programme

AGE	Days	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
TEMPERATURE	Under the heaters	35–36				32–34			28–30							
	Living area	28					27–28			25–26						
	Whole house heating	31–32	30–31	29–30	28–29		26–27									
HUMIDITY	%	50–60														
EQUIPMENT	Drinker	1 round drinker for 60 birds; 1 nipple for 8 birds								1 round drinker for 80 birds; 1 nipple for 8–10 birds						
	Feeder	1 feeder/50–70 birds + chick paper (> 50%)								Chain feeder: 6–8 cm/bird; 1 oval pan feeder/14–16 birds; 1 round pan feeder/13–15 birds						
DENSITY	Chicks/m ²	25–30					12			7–10						
LIGHTING PROGRAMME	Light duration (h)	24	22		20		18			16						
	Light intensity (lux)	80 (If permitted by local regulations)						60 (If permitted by local regulations)								
FEED	Ad libitum	Ad libitum for 3 weeks and until 4 weeks if the 3 weeks bodyweight is not achieved														
	Feed type	Pre–starter crumbles or Mini pellets														

Information icons and QR codes linking to Nutrition Recommendations and Hubbard Poster Brooding PS.

3. REARING PERIOD (0–20 WEEKS): HOW TO CHOOSE THE OPTIMUM LIGHTING PROGRAMME? DARK PRODUCTION HOUSE

Numerous parameters can affect sexual maturity including: house type, latitude, season, bodyweight curve and uniformity pattern. On page 7 and 8 some examples are provided the general principles to design a lighting programme for each female breed. Please consult your local Hubbard Technical Specialist, especially if males have to be housed in the same house as females.

► REQUIREMENTS FOR AN OPTIMAL LIGHTING PROGRAMME



► Rearing conventional males (M77 / M99) with females in the same rearing house is not recommended → See “*PS males mated with Premium females management guide*”.

- For solid-sided houses use light traps on fans and all air inlets (including the minimum ventilation).
- For side-curtain tunnel housing it is advised to darken the houses for the whole life of the flock, using solid black curtains.
- For open-sided traditional houses take care if using dew/shed net to cover the sides at all ages. Ventilation and season should be carefully considered to ensure optimum flock and litter conditions to maintain good footpad condition.



Open-sided rearing house with black curtains



Dark rearing house – light traps on fans and inlets



Dark rearing house with natural light leakage




Dark rearing house – no natural light leakage

Case N°1: DARK REARING HOUSE / DARK PRODUCTION HOUSE

TABLE 1: LIGHTING PROGRAMME ACCORDING TO THE BREED

REARING HOUSE	DARK		
PRODUCTION HOUSE	DARK		
Breed	JA57 – JA57Ki – P6N	JA87 – REDBRO MINI	Light intensity *
Flock age	Light duration *		
14–20 d	16 h	16 h	40 lux
21–28 d	16 h	15 h 30	10 lux
29–35 d	15 h 30	15 h	
36–42 d	15 h	14 h	
43–49 d	14 h 30	13 h	
50–56 d	14 h	12 h	
57–63 d	13 h	11 h	
64–70 d	12 h	10 h	
71–77 d	11 h	10 h	
78 d to transfer (126–140 d)	10 h	10 h	

* Hours of light and light intensity may both be subject to local regulations.



Light duration for Premium Mini PS females–Dark production houses

3. REARING PERIOD (0–20 WEEKS): HOW TO CHOOSE THE OPTIMUM LIGHTING PROGRAMME? OPEN-SIDED PRODUCTION HOUSE

Case N°2: DARK REARING HOUSE / OPEN-SIDED PRODUCTION HOUSE

TABLE 2: LIGHTING PROGRAMME ACCORDING TO THE BREED


REARING HOUSE	DARK			
PRODUCTION HOUSE	OPEN-SIDED			
Breed	JA57 – JA57Ki – P6N		JA87 – REDBRO MINI	
Natural daylength at 21 weeks of age	< 10 h	> 10 h	< 10 h	> 10 h
Flock age	Light duration *			
14–20 d	16 h	16 h	16 h	16 h
21–28 d	16 h	16 h	15 h 30	16 h
29–35 d	15 h 30	15 h 30	15 h	15 h 30
36–42 d	15h	15h	14 h	15 h
43–49 d	14 h 30	14 h 30	13 h	14 h 30
50–56 d	14 h	14 h	12 h	14 h
57–63 d	13 h	13 h 30	11 h	13 h
64–70 d	12 h	13 h	10 h	12 h
71–77 d	11 h	12 h 30	10 h	12 h
78 d to transfer	10 h	12 h **	10 h	12 h **
Transfer age	140 d	147 d	140 d	147 d

* Hours of light and light intensity may both be subject to local regulations.


** In some cases (too early mature females in rear), the 12 h plateau could be reached at 14 weeks of age.

LIGHT INTENSITY MANAGEMENT


- It is essential to provide uniform light at bird level in each pen all along the rearing period.
- The use of an appropriate light meter at bird level brings precision to set the light bulbs at the correct distance from the floor.
- Replace the light source as soon as possible when a bulb is not working properly.
- Remove the dust from the light source when required.



Dark rearing house – good light distribution



Light duration for JA57 - JA57Ki - P6N - Open-sided production houses



Light duration for JA87 - REDBRO MINI - Open-sided production houses

3. REARING PERIOD (0–20 WEEKS): HOW TO FOLLOW THE BODYWEIGHT TARGETS?

MANUAL WEIGHING



Every week, always the same day: hatch day is the best.




Best before feeding.

After feeding, use a with feed and water bodyweight target.




Age < 2 weeks

Age > 2 weeks



Check calibration

Check calibration



- Take samples from 3 separate places but not too close to the main feed hopper. Always keep the same location. **Sample size: 3–5% of the flock size or at least 100 birds per pen.**
- Weigh all the birds of the catching pen.

AUTOMATIC WEIGHING




Every week, always the same day: hatch day is the best.



Best before feeding.

After feeding, use a with feed and water bodyweight target.

Check calibration




Results from an automatic scale

- Ensure to place the scales as soon as possible to maximize the number of birds weighed.
- The lower and upper limits of the bodyweights often need to be adjusted.
- If an unexpected variation of the bodyweight from the objective is observed, it is recommended to re-weigh by hand a new sample.



Analysis of the weekly results for an optimal feed allocation management:

1. The feed allocation per bird should be based on the average body weight calculated after weighing.
 2. Compare the average bodyweight with the bodyweight target. Plot the average weight on the growth curve to see the trend.
 3. Calculate the weekly gain and compare it with the objective mentioned in the Performance Objectives of the female breed used.
 4. Adjust feed allocation according to the weekly weight gain achieved and the one targeted for the next week. Please do not strictly follow the feed intake targets as they are only provided as a guideline and consider actual feed increases.
- Be prepared to control feed allocation from 8 weeks of age to avoid excess bodyweight gain after 10 weeks of age.
 - Transition feed (Pre-lay feed) is not recommended diet as it may promote growth and too early sexual maturity. → See "**Nutrition Recommendations**".
 - If the first eggs are laid before 140 days, please contact your Hubbard Technical Specialist.



3. REARING PERIOD (0–20 WEEKS): HOW TO REACH AND MAINTAIN GOOD FLOCK UNIFORMITY AND GOOD APPETITE?

There are two possible indicators to express flock uniformity:

1. The uniformity is defined by the percentage of birds weighing between $\pm 10\%$ of the average bodyweight of the flock/pen. → See chart below.
2. The coefficient of Variation (CV) is the variation of bodyweights within the flock (= Standard deviation/average flock bodyweight). The lower the CV%, the more uniform the flock is.

GRADING: OPTIONAL

- 7 days: grade or at least isolate the smallest chicks by eye and manage them with special care to achieve target weight by 4 weeks.
- 21–28 days: if needed grade 100% of the flock and create weight groups, each with at least 85% uniformity (CV < 7.5). → See Bulletin *“Grading of broiler breeders”*.
- 29–84 days: aim to maintain or improve flock uniformity by having a close follow up of the feed distribution. → See Poster *“Quality of feed distribution”*.
- 84–98 days: if flock uniformity is less than 75% (CV > 9), re-grade the flock to assure that each weight group achieves at least 85% uniformity (CV < 7.5). Have a close bodyweight follow up during the vaccination periods. Anticipate this with extra feed.

FEED PROGRAMME

- Daily feeding gives successful results in many situations.
- If fractionated feeding is considered then please contact your Hubbard Technical Specialist to discuss possible solutions to optimise feed distribution.
- Adjust the feed depth in the feeders each time the feed allocation is changed to assure the correct distribution.
- Use of a low energy grower feed (< 2 650 kcal/kg) is preferred to improve gut health and eating behaviour. If high fibre sources are added to the feed these must be of consistent quality and mycotoxin free. → See Bulletin *“Dietary fibre”*.



SPIN FEEDING

- Spin feeders are preferably used from 35 days of age when females are large enough to consume pellets (1 spin feeder per pen). If introduced earlier then special care must be taken to ensure small females are able to consume feed.
 - ▷ Check growth is not depressed and ensure the target weight at 42 days is achieved.
 - ▷ Consider extending the use of feed provided on paper or feeders to complement spin feeders.
- Avoid pellets over 3.2 mm diameter where possible and cut pellets as short as possible while keeping pellet quality. Ensure small chicks can consume the pellet.
- At the end of the rearing period when the birds are transferred to production house, please pay special attention to ensure the birds find their new feeders and drinkers. This may require additional temporary drinkers to be placed on the floor or slats.

i




[Hubbard Bulletin
Dietary fibre](#)



[Hubbard Bulletin
Grading](#)



[Hubbard Poster
Feed distribution](#)



Results of one manual weighing – % uniformity definition

3. REARING PERIOD (0–20 WEEKS): HOW TO PROMOTE GOOD BEHAVIOUR AND PROVIDE GOOD ANIMAL WELFARE?

For all enrichments it is important to consider the risks to compromise biosecurity, especially from straw or other bales and material brought into the house and also the ease of washing of fixed enrichments like platforms.

► PERCHES AND PLATFORMS

From 28 days, the use of a perching system is strongly recommended to stimulate activity and to train birds to jump onto the slats and manual nests. This helps prevent floor eggs.

- If the production house is equipped with automatic nests:
 1. Provide 4 m² of platform for 1 000 birds.
 2. Optimal position of the platform is under the drinking system (see picture below).
- If the production house is equipped with manual nests:
 1. Provide 5 cm of perch space per bird.
 2. Place some manual nests from 12–14 weeks of age.
- Platforms and perches should not have sharp edges that may damage birds.




Platforms



Perches

► WATER MANAGEMENT/WATER QUALITY

- With a regular and proper management of drinker height and water flow according to flock age, no controlled access to water should be needed.



Correct bell drinker height

Correct nipple height

- Regularly check the chemical and bacteriological water quality to ensure that water sanitation is working properly.

1. Optimal water pH: 5.5 to 6.8.
2. Optimal ORP higher than 600 mV with chlorine.
3. Chlorine concentration at the end of the pipe: target = 1 ppm (acceptable range: 0.5 to 3 ppm).



► GRIT AND GRAIN: optional.

- Insoluble grit to promote the gizzard development.


1. ø 2–3 mm: 3–5 g/bird/week from 5 weeks of age.
2. ø 3–5 mm: 3–5 g/bird/week from 10 weeks of age.




- Sanitized scratch grain (cracked maize or whole wheat): 3 g/bird/day 4–5 h after feeding from 5 weeks of age.

► ENRICHMENT: optional unless required by local regulations.

- White ropes – max 20 cm long – no contact with the litter.
- Wood shaving balls – 1 piece for 500 to 1 000 birds, placed on floor.
- Pecking blocks – 1 piece for 500 to 1 000 birds – Consider the hardness of the material.
- Alfalfa balls – 1 piece for 500 to 1 000 birds, hanged or placed on floor.



Example of white ropes



Example of pecking blocks



Example of pecking blocks



Example of alfalfa balls

4. PRODUCTION PERIOD (20–65 WEEKS): HOW TO MAKE THE FIRST 10 DAYS A SUCCESS? FOCUS ON LIGHT

The first 10 days after the first egg are very important to achieve good results in terms of total eggs, hatching eggs and nest eggs. Eating behaviour, stocking density, eating/drinking space and ventilation need to be managed properly for a good peak and persistency of lay.

LIGHTING PROGRAMME: the objective is to reach 5% weekly production at 22 weeks of age (JA57, JA57Ki) and 23 weeks of age (JA87, Redbro Mini, P6N).

► Light duration:

1. Refer to the Table 3 and 4 to see some examples of lighting programme. Be prepared as the onset of lay should normally start about 1–2 weeks after the first light stimulation.
2. Provide the maximum light intensity and 14 h light by 20% daily production without ever exceeding 16 h light duration.
3. For open-sided houses: please contact your Hubbard Technical Specialist to define how to provide the additional light.

► Light intensity:

1. Do not reduce the light intensity between the rearing house and the production house.

Ensure bright light (60–80 lux) in the mating area and a lower light intensity level on the nests.

2. Rapid increase of light intensity is important with pan feeders so females see feed clearly before feed intake is increased with production.
3. Ensure that the light distribution is uniform especially during the night time for open-sided houses. If possible, it is better to simulate dawn in the morning and dusk in the evening. This may be required under local regulations.

► Lighting systems:

1. Make the best choice of light type such as incandescent, energy saver (white or yellow), fluorescent tube, sodium and LED to assure uniform light intensity at bird level.
2. Good results are usually achieved with typical color temperatures between 2 700–4 000 K. There is no clear evidence that color temperatures above 4 000 K give better results.



Case N°1: DARK REARING HOUSE/DARK PRODUCTION HOUSE

TABLE 3: LIGHTING PROGRAMME ACCORDING TO THE BREED

REARING HOUSE	DARK		
PRODUCTION HOUSE	DARK		
Breed	JA57 – JA57Ki	JA87 – REDBRO MINI – P6N	Light intensity *
Flock age (d)	Light duration *		
Transfer age	10 h at 126–140 d	10 h at 126–140 d	10 lux
148 d	12 h	10 h	20 lux
154 d	14 h	12 h	40 lux
161 d	14 h or 15 h **	14 h	60–80 lux
From 168 d	14 h or 15 h **	15 h **	60–80 lux

* Hours of light and light intensity may both be subject to local regulations.

**Provide 15 h light if some floor eggs are observed just before the lights come on.

Case N°2: DARK REARING HOUSE/OPEN-SIDED PRODUCTION HOUSE

TABLE 4: LIGHTING PROGRAMME ACCORDING TO THE BREED

REARING HOUSE	DARK				
PRODUCTION HOUSE	OPEN-SIDED				
Breed	JA57 – JA57Ki		JA87 – REDBRO MINI – P6N		Light intensity *
Natural daylength at 21 weeks of age	< 10 h	> 10 h	< 10 h	> 10 h	
Flock age (d)	Light duration *				
Transfer age	10 h at 140 d	12 h at 147 d	10 h at 140 d	12 h at 147 d	10 lux
148 d	12 h	14 h	10 h	12 h	Natural daylight +artificial light (60–80 lux)
154 d	14 h	15 h	12 h	14 h	
161 d	15 h **	15 h	14 h	15 h **	
From 168 d	15 h **	15 h	15 h **	15 h **	

* Hours of light and light intensity may both be subject to local regulations.

**Provide 15 h light if some floor eggs are observed just before the lights come on.


4. PRODUCTION PERIOD (20–65 WEEKS): HOW TO MAKE THE FIRST 10 DAYS A SUCCESS? FOCUS ON FEED PROGRAMME

FEED PROGRAMME: Maximising appetite is crucial to ensure a good early egg size and maximising the nest eggs during onset of lay.

- Flocks transferred from the rearing farm could lose bodyweight.
 - Anticipate this with extra feed especially during the vaccination periods and when the feeding system is different in the production house.
 - Give consistent feed increases until the flock reaches 5% daily egg production.
- Increase feed according to the production level:
 - Once 5–10 % daily production, feed the flock according to the daily increase in production: typically + 4–5 g/day. Typical increase in feed is 0.8–1 g per 1% increase in daily egg production. Reach the maximum feed intake by 55–60% daily production. → Refer to the relevant Premium Breed *“Performance Objectives”*.
 - The onset of lay may be very rapid and it is advised to be well prepared. For example, it can take only 12–14 days from 5% to 80% daily production.
 - Ensure the daily egg weight increase is consistent using the daily onset of lay sheet. The Poster *“PS Premium Mini Egg weight management”* explains the egg weighing method.
 - Monitor feed clearance time at peak feed on a daily basis (around 8–10 hours for JA57, JA57Ki and P6N, 6–8 hours for JA87, Redbro Mini) but may vary according to feed and environment.
 - Ensure feeders are empty for at least 3 hours every day, preferably in the middle of the day (around 5–6 hours after light on) as shown in the pictures below. If this is not the case:
 1. Consider reducing the total amount of feed distributed per day to ensure this empty period is observed during some days.
 2. Do not add extra feed distributions during the day during hot weather.
- Risk factors for low appetite or long feed clearance time:
 - Excess dietary protein may cause longer feed clearance time and then it will be difficult to manage the balance between nutrient supply for egg production and body maintenance.
 - During hot weather (house temperature > 28 °C) to stimulate production and reduce heat stress:
 1. Consider the use of the “onset of lay” feed formula or hot climate feed formulation (higher energy feed to reduce total volume) → See *“Premium Mini PS Nutrition Recommendations”*.
 2. Maximise the physical quality of the feed or increase the particle size of the feed (mash to crumble or crumble to pellet).
 3. Turn on the lights in the middle of the sleeping period for 1 to 2 hours and give cold water with vitamin C (if permitted by local regulations).

EXAMPLE OF A GOOD ONSET OF LAY FOR A PREMIUM MINI PS FLOCK

- The stimulation of appetite early in production determines the consumption capacity at peak a few days later as shown in the chart below.
- If the daily feed increases have not kept up with the production, the peak feed level and likely also production will be lower.



* It is important that feed level stays ahead of the of the egg production curve.



Different levels of empty feeders with chain feeders

i

[Hubbard Poster
JA57 & JA57Ki
Egg weight management](#)

[Hubbard Poster
JA87 & REDBRO MINI
Egg weight management](#)

[Hubbard Poster
P6N
Egg weight management](#)

4. PRODUCTION PERIOD (20–65 WEEKS): HOW TO MAKE THE FIRST 10 DAYS A SUCCESS? FOCUS ON FEED DISTRIBUTION

QUALITY OF FEED DISTRIBUTION

► Ensure on a daily basis that feed remains evenly distributed into the feed system to maintain a good flock performance (flock uniformity, egg production, egg weight).

▷ Ensure a uniform feed level between the feeder lines. Some litter can accumulate and block the feed outlet.

▷ Adjust the time of the last feed turn (see figure below).

▷ Fast feed distribution (< 4 minutes).



▷ Maintain the supplementary feed hoppers until the end of the flock.

▷ Management of pan feeders: distributing the daily feed in one or two meals may require careful attention, particularly by maximising the first feed distribution. If this cannot be achieved:

1. Add an extra feed distribution, but avoid feeding during the next 5–6 hours, which coincides with the daily peak of egg production.

2. Keep the feed depth at the maximum setting after transfer so that females can easily see and access the feed in the pans.

► Different methods for the feed distribution: whatever the method chosen the number of times per day feed is distributed should be kept to a minimum (< 4 feed turns).

▷ Early feeding method: the first feed turn is given within 30 minutes after lights on.

1. After transfer, the birds can be fed only in the morning if the feed clearance time is not longer than 6–7 hours.

2. If it takes longer, it is recommended to implement two meals during the day (e.g. 60% in the morning and 40% at least 6 hours after the first feed distribution, to be adjusted according to flock appetite).

▷ Late feeding method: the first feed turn is given a few hours after the laying period like for example: 70% of the daily feed amount at noon and 30% in the late afternoon.

▷ Night feeding method: when the flock is raised in hot climate, it is possible to use a night feeding programme (supplementary feed):

1. Applying 2 hours of light starting 2 hours after lights off (resting period).
2. Feeding times could be adapted depending on the local conditions.



TABLE 5: ADVANTAGES AND DRAWBACKS OF THE FEED DISTRIBUTIONS METHODS

METHOD	ADVANTAGES	DRAWBACKS
Early feeding	► Securing egg shell quality ► Good flock activity in the afternoon	► Non optimal nesting behavior under challenging conditions
Late feeding	► Easier to check the quality of the feed distribution ► Improved nesting behavior	► Not recommended in hot climate.
Night feeding	► Securing the feed intake in hot climate.	► Disruption of the egg-laying window.

4. PRODUCTION PERIOD (20–65 WEEKS): HOW TO MAKE THE FIRST 10 DAYS A SUCCESS? FOCUS ON NEST LAYING

GENERAL RECOMMENDATIONS FOR NEST LAYING

- Transfer birds directly on slats and provide water and feed (if feeders are on the slats).
- According to the type of males being used, mixing to ensure 7.5–9.5% males of the appropriate sexual maturity at 20–22 weeks. More than 9.5% males could be risky to maintain a low floor eggs %.
→ See “*PS males mated with Premium females management guide*”.
- Collect floor eggs at least every hour during the laying time. From very first egg, this may be 10–12 times per day.
- Provide a low level of litter after transfer (1–3 cm in hot conditions/4–6 cm in cooler conditions). In case of using less than 3 cm, add fresh litter on a regular basis to maintain good litter quality.
 - Place dummy eggs or marked eggs in the nests to attract birds.
 - Avoid draughts of cold air or very high air speed inside the nests.
- Do not feed at the time when most eggs are laid during the day.
- A full checklist on the nesting behavior is available from your local Hubbard Technical Specialist.

i



[Male management guide](#)

SPECIFICATIONS FOR AUTOMATIC NEST


- Open the nest during the day at least 2 weeks before first egg. It is also possible to open nest flaps until 30% daily production.
- Turn-on lights inside nests one hour before lights on in the house to help early laying birds find their way to the nests.
- Nest belts can be turned-on twice during the day to familiarise birds with the noise.
- If slat height > 40 cm, place steps to help birds jump onto the slats. Slat slope should not exceed 5–8°.
- Ensure a correct nest ratio (80–90 females/linear meter for colony nests and 4 females per nest for individual nest systems).
- Adequate nipple flow (70–100 ml/min) and a correct distance (> 1 m) from the nest to avoid a fence effect in front of the entrance to the nests.
- Using bricks on the slats may help birds distribute more evenly in the nests.



Slat steps to promote nest access




Bricks on the slats close to the corners



Nest flaps opened until 30% daily lay

SPECIFICATIONS FOR MANUAL NEST

- Introduce nests at least 4 weeks before the onset of lay.
- Put clean bedding material in the nests to make them attractive.
- Perches must be large enough to allow easy access to both the 1st and 2nd nest level.
- Ensure the correct nest box ratio (4 females/nest).
- If egg collections are made at a time when females are laying most of the eggs, avoid disturbing females in the nests.
- Lower the nests close to the litter level to promote nest access.



Good perch design for an easy nest access

4. PRODUCTION PERIOD (20–65 WEEKS): HOW TO MAXIMISE THE PERSISTENCY OF LAY?

Objectives in order to maximise the egg production and the female liveability:

1. Control flock bodyweight, flock uniformity, and body fat.
2. Ensure egg weight is on target.

► FEEDING MANAGEMENT

- Check the evolution of the egg weight and female growth before reducing the feed intake. If production drops after a feed reduction, reinstate the previous amount.
- Adjust feed allocation for both hot (> 28°C) and cold (< 18°C) weather to meet metabolic requirements.

► FEED FORMULA

- It is recommended to change to Breeder II diet once the egg weight reaches 58 g (hot climates/hot season) or 60 g (cold climates/cold season).
- Check the physical quality of the feed on a regular basis using the Hubbard feed sieve and related Hubbard calculation software. → See “*Hubbard Feed Sieve Tool*”.

► FEMALE CONDITION/BODYWEIGHT CONTROL

- Weekly control of female bodyweight is essential and feed should be immediately adjusted at any time that bodyweight drifts away from the recommended objective.
- Assess abdominal fat on a regular basis. Fatness should be controlled using Breeder II feed and timely management of the feed allocation.
- Assess female feathering on a regular basis.



► LIGHTING PROGRAMME

- It is possible to increase an extra hour after 40 weeks of age, but without exceeding 16 h light duration.

► WATER QUALITY: regularly check the chemical and bacteriological water quality to ensure that water sanitation is working properly.

- Ideal water pH: 5.5 to 6.8.
- Optimal ORP or Redox potential higher than 600 mV with chlorine (450 mV with chlorine dioxide and 300 mV with peracetic acid or peroxide).
- Chlorination concentration at the end of the pipe: target = 1 ppm (Acceptable range: 0.5 to 3 ppm).

► BROODINESS: Below is a list of the risk factors.

- Feed problems: formulation, presentation or feed distribution combined with insufficient feed intake before and immediately after peak egg production (low nutrient particularly energy intake).
- Bird environment: high environmental temperature inside the house and the nests.
- Nest management: infrequent egg collections and visible belt eggs.
- Poor management of floor eggs.
- Insufficient water consumption.
- Viral diseases and/or parasites.
- Make daily checks for the presence of non-laying birds from the peak of lay onwards. Isolate broody hens away from nests in a pen for one week.





[Hubbard Feed Sieve Tool](#)



[Hubbard Poster JA57 & JA57Ki Egg weight management](#)



[Hubbard Poster JA87 & REDBRO MINI Egg weight management](#)



[Hubbard Poster P6N Egg weight management](#)



[Hubbard Bulletin Water quality](#)

5. BIOSECURITY

LITTER MANAGEMENT



Litter/bedding storage in an adapted biosecure area to maintain the pathogen-free status (inside the house).



No accumulated litter at the end of the cycle.


SHED TO SHED ACCESS



If corridor between the two houses provide three-zone entry.



If no corridor between the houses, provide one shower unit in each house.



VISITORS



Restrict visitors and adapt quarantine period.



Provide shower facilities.



Provide change of clothes and boots.

VEHICLES



Vehicles must be cleaned prior to entering the farm facility.



DO NOT allow unauthorised vehicles on the farm.



Make sure all feed and gas deliveries stay outside of the fence.



Shoe covers are to be worn on shoes from cars into the shower facilities.

EXTERIOR



Do not treat the concrete or area inside the fence as clean.

INTERIOR



Easy to clean and disinfect concrete floor.



Easy to clean and disinfect equipment. The resting period ideally lasts for at least 10 days.



Ensure good quality water is available.

ANIMAL CONTROL



Keep other livestock away from facility.

PEST CONTROL




Establish regular baiting for rodents and insects, prevent access of wild birds according to local regulations.

KEEP A RECORD



Visitors, feed/litter delivery, cleaning and disinfecting, pest control.



- 1 Freezer room: disposal of dead birds in a biosecure freezer.
 - 2 Gas tank
 - 3 Fumigation room
 - 4 Egg storage room
 - 5 Vehicle disinfection unit
 - 6 Staff room
- ➡ Go forward principle

* It may be possible to rear different Hubbard breeds in the same house. Please consult your local Hubbard Technical Specialist.

Handwriting practice lines consisting of 20 horizontal dotted lines.

Lined area for notes.





The performance data contained in this document was obtained from results and experience from our own research flocks and flocks of our customers. In no way does the data contained in this document constitute a warranty or guarantee of the same performance under different conditions of nutrition, density or physical or biological environment. In particular (but without limitation of foregoing) we do not grant any warranties regarding the fitness for purpose, performance, use, nature or quality of the flocks, nor any warranty regarding compliance with local legislation regarding health, welfare, or other aspects of animal production. Hubbard makes no representation as to the accuracy or completeness of the information contained in this document.



AMERICAS
HUBBARD LLC
123 Gallus Rd
PIKEVILLE, TN 37367 – U.S.A.
TEL. +1 (423) 447 6224
contact.americas@hubbardbreedersusa.com

EUROPE, MIDDLE EAST, AFRICA
HUBBARD S.A.S.
Mauguérand
22800 LE FOEIL – FRANCE
TEL. +33 2 96 79 63 70
contact.emea@hubbardbreeders.com

ASIA
HUBBARD S.A.S.
Mauguérand
22800 LE FOEIL – FRANCE
TEL. +33 2 96 79 63 70
contact.asia@hubbardbreeders.com

Hubbard is a registered trademark of Hubbard in the US and other countries.
All other brands and trademarks are the trademarks of their respective owners.

© 2026 Hubbard